



Contraindicated Drugs in Parkinson Disease Patients



WISCONSIN
PARKINSON
ASSOCIATION

16655 W. Bluemound Road, Suite 330
Brookfield, WI 53005

414-312-6990 | wiparkinson.org

8/18

The drugs listed on this card have the potential to cause or worsen the symptoms of Parkinson disease and should not be prescribed for an individual with a Parkinson's diagnosis.
Always ask your doctor before starting or discontinuing any medication.

	Brand Name	Chemical Name
Antipsychotic (used for agitated confusion)	Haldol	Haloperidol
	Loxitane	Loxapine HCL
	Mellaril (high dose)	Thioridazine
	Moban	Molindone
	Navane	Thiothixene
	Orap	Pimozide
	Prolixin, Permitil	Fluphenazine
	Serentil	Mesoridazine Besylate
	Stelazine	Trifluoperazine
	Taractan	Chlorprothixene
	Thorazine	Chlorpromazine
	Trilafon	Perphenazine
Antidepressant	Ascendin	Amoxapine
	Marplan	Isocarboxazid
	Nardil	Phenelzine
	Parnate	Tranlycypromine
	Triavil	Combination of Perphenazine & Amitriptyline
Antivomiting	Compazine	Prochlorperazine
	Inapsine	Droperidol
	Phenergan	Promethazine
	Reglan	Metoclopramide
	Torecan	Thiethylperazine
Miscellaneous, blood pressure, post- op medication	Moderil	Rescinnamine
	Rauverid	None
	Rauwiloid	None
	Serpasil	Reserpine
	Wolfina	None
Avoid unless benefit outweighs risk	Abilify	Aripiprazole
	Aldomet	Alpha-methyldopa
	Depakote	Divalproex Sodium
	Eskalith, Lithobid	Lithium
	Geodon	Ziprasidone
	Pavabid	Papaverine
	Risperdal	Risperidone
	Zyprexa	Olanzapine
Narcotic analgesic	Demerol*	Meperidine
Cough suppressant	Numerous brands*	Dextromethorphan
Decongestants	Sudafed*	Pseudoephedrine
	Sudafed PE*	Phenylephrine
		Ephedrine
	<i>*should not be used with Azilect (Rasagiline) or Eldepryl (Selegiline)</i>	